



**West African Ornithological Society**  
**Société d'Ornithologie de l'Ouest**  
**Africain**



Join the WAOS and support  
the future availability of free  
pdfs on this website.

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.  
If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

Devenez membre de la  
SOOA et soutenez la  
disponibilité future des pdfs  
gratuits sur ce site.

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.  
Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante  
(2) pour économiser du papier.

There are no real surprises in the inclusions and absences in the species listed, apart from Malimbus cassini. Sight records of males (only) of this species from Tafo make me recall being invited there in 1960 by Gordon Donald because he was convinced he had sighted M. ibadanensis on the grounds of the West African Cocoa Research Institute at Tafo. In those days WACRI had research stations at both Tafo (Ghana) and Ibadan (Nigeria) and Donald's duties caused him to move between the two places, so he knew ibadanensis well from field sightings and hand specimens from around Ibadan. He was sufficiently convinced of its occurrence at Tafo to get me there to look with him, but no sighting was obtained during my all too brief stay. It is noteworthy that both Hall & Moreau (1970 Atlas of Speciation in African Passerine Birds, BM (NH), London) and Louette (1981, The Birds of Cameroon, Kon. Acad. Wetensch. Lett. Schone Kunst. België, Brussels) state that the Sanaga River in Cameroon is the western limit of M. cassini. From an inspection of skins of both species at the British Museum it would seem that male cassini has the red throat/breast intermediate in extent between the red 'bib' of female ibadanensis and the red 'apron' of the male, making field distinction of the two species very difficult if not impossible. The entirely black female cassini is also easily confused with other black weavers. There must be a case for collecting at least one specimen to resolve which species actually occurs in Ghana.

Since a Table (11, p.51) analyses the differences between the avifaunas of Ghana and Nigeria, a feature I found of special interest, I thought it a pity that neither the species by species treatment nor the Status Table had a symbol to indicate absence from Nigeria. The comparison of the two avifaunas would have been made more complete by a list of the species found in Nigeria but so far not reported from Ghana.

The Ghana Checklist rightly draws attention to the relative paucity of breeding data and suggests that this should be a field for future workers to investigate. There are a surprising number of species for which there are no Ghana breeding records but with breeding noted elsewhere in West Africa, notably Nigeria. This probably reflects the relative numbers of observers in the different countries.

Finally the Bibliography warrants favourable mention and the Gazetteer accurately pinpoints the places mentioned.

J.H. Elgood

BIRDS OF THE EASTERN CAPRIVI by J.H. Koen (1988) Southern Birds 15: 1-73 ISBN 0-620-10490-2. SA R7.00, from PO Box 650284, Benmore 2010, South Africa.

The avifauna of the end of the long finger-like projection of Namibia sandwiched between Botswana and Zambia. Includes an annotated checklist of 374 bird species and a description of geography, climate and habitats.

H.Q.P. Crick