

West African Ornithological Society Société d'Ornithologie de l'Ouest Africain



Join the WAOS and support the future availability of free pdfs on this website.

http://malimbus.free.fr/member.htm

If this link does not work, please copy it to your browser and try again. If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

Devenez membre de la SOOA et soutenez la disponibilité future des pdfs gratuits sur ce site.

http://malimbus.free.fr/adhesion.htm

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer. Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante (2) pour économiser du papier.

- DOWSETT, R.J. &FORBES-WATSON, A.D. (1993) Checklist of Birds of the Afrotropical and Malagasy Regions. Tauraco Press, Liege.
- HANCOCK, J.A., KUSHLAN, J.A. & KAHL, M.P. (1992) Storks, Ibises and Spoonbills of the World. Academic Press, London.
- SALEWSKI, V., BOBEK, M., PEŠKE, I. & POJER, F. (2000) Status of the Black Stork *Ciconia nigra* in Ivory Coast. *Malimbus* 22: 92–93.
- SNOW, D.W. & PERRINS, C.M. (1998) The Birds of the Western Palearctic, Concise Ed., vol. 1. Oxford University Press, Oxford.
- THIBAULT, J.-C. & BONACCORSI, G. (1999) *The Birds of Corsica*. Checklist 17, British Ornithologists' Union, Tring.
- WALSH, J.F. (1991) On the occurrence of the Black Stork Ciconia nigra in West Africa. Bull. Brit. Orn. Club 111: 36-42.

Received 7 April 2001

Revised 1 July 2001

J. Frank Walsh

80 Arundel Road, Lytham St. Anne's, Lancashire FY8 1BN, England

First breeding record of Little Grey Woodpecker *Dendropicos elachus* in Nigeria

On 11 Jan 2002, in relatively dense Acacia woodland 2 km east of Alagarno village (13°09'N, 13°33'E), Borno State, Nigeria, I noticed a male Little Grey Woodpecker Dendropicos elachus repeatedly chasing a pair of Bush Petronia Petronia dentata from a 6-m Acacia tortillis. The behaviour strongly suggested nesting and I located a 5 cm diameter entrance hole at 1.5 m height, on the underside of a 10 cm diameter dead branch. It was impossible to look into the cavity but, on moving a short distance from the tree, the woodpecker reappeared with a small food item in its bill. On calling, a female left the cavity and flew off. The male entered the cavity briefly, reappeared without the food and flew off. There are no previous records of this species breeding in the country (Elgood et al. 1994) and few nests have been described throughout the species' range (Fry et al. 1988).

Little Grey Woodpecker is an uncommon resident of the Sahel zone from Senegambia east to Sudan (Fry et al. 1988). In Nigeria it is restricted to the extreme NE, near Lake Chad. The habitat of this nest is typical of the species in Nigeria, though it also occurs in more open *Acacia* and mixed *Acacia-Balanites* woodland in the Alagarno area (pers. obs.).

References

BARLOW, C., WACHER, T. & DISLEY, D. (1997) A Field Guide to the Birds of The Gambia and Senegal. Pica Press, Mountfield.

BORROW, N. & DEMEY, R. (2001) Birds of Western Africa. Helm, London.

ELGOOD, J.H., HEIGHAM, J.B., MOORE, A.M., NASON, A.M., SHARLAND, R.E. & SKINNER, N.J. (1994) *The Birds of Nigeria*. Checklist 4 (2nd ed.), British Ornithologists' Union, Tring.

FRY, C.H., KEITH, S. & URBAN, E.K. (1988) The Birds of Africa, vol. 3. Academic Press, London.

Received 9 February 2002

Jared M. Wilson

A.P. Leventis Ornithological Research Institute, Jos, Nigeria and Edward Grey Institute, Dept of Zoology, South Parks Road, Oxford, OX1 3PS, U.K. e-mail: jared.wilson@zoo.ox.ac.uk