

West African Ornithological Society Société d'Ornithologie de l'Ouest Africain



Join the WAOS and support the future availability of free pdfs on this website.

http://malimbus.free.fr/member.htm

If this link does not work, please copy it to your browser and try again. If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

Devenez membre de la SOOA et soutenez la disponibilité future des pdfs gratuits sur ce site.

http://malimbus.free.fr/adhesion.htm

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer. Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante (2) pour économiser du papier.

Good, A.I. (1952) The birds of French Cameroon - Part 1. Mem. Inst. fr. Afr. noire Cent. Ca eroun No 2.

Good, A.I. (1953) The birds of French Cameroun Part 2. Mem. Inst. fr. Afr. noire Cent. Cameroun No 1

Reis, J.A. (1945) Les oiseaux du Cameroun française. Bull. Soc. Etud. Cameroun 11, 17-55.

Walker, R.B. (1965) Sokoto birds: Addendum. Bull. Niger. Om. Soc 1(4), 20.

----000000----

THE DESTRIBUTION OF THE RED-EYED TURTLE DOVE STREPTOFELIA SEMITORQUATA

This note raises a query regarding the habitat of the Red-eyed Turtle Dove which can be answered by members of the Society. The query arose when I was writing up notes on the birds seen on a short visit to Mount Cameroun in December 1970. (see this issue)

In my experience in Ghana it is not found in the lowland forest areas and only occurs in the more open and drier areas to the north of the forest belt and on certain areas of the Acera Flain. This distribution agrees with Bannerman (1953) who writes that it is "a bird of the open country and not of the forest, preferring plantations and second growth". In Cameroun, however, I found the bird to my surprise in the montane forest and Serle (1965) writes that the bird is found "in the forest from sea level up to 6000 ft. and in the orchard bush and gallary woods of the savanna." This distribution in Cameroun agrees with Machworth-praed and Grant (1971) who write that it is "found throughout the area (West Africa) in wooded country or forest, but quite often seen also in open country..." Presumably in Nigeria there is a transition zone and it would be of interest if members would write in to the Bulletin describing the habitat in which they have found the Red-eyed Turtle Dove.

REFERENCES

Bannerman, D.A. 1953. The birds of West and Equatorial Africa. 2 Vols. Oliver and Boyd.

Machworth-Praed, C.W. and Grant, C.H.B., 1971 Birds of West Central and Western Africa, Longmans, London.

Serle, W., 1965. A third contribution to the ornithology of the British Cameroun. Ibis 107: 60-94, 230-246.

L. Grimes.