



**West African Ornithological Society  
Société d'Ornithologie de l'Ouest  
Africain**



**Join the WAOS and support  
the future availability of free  
pdfs on this website.**

<http://malimbus.free.fr/member.htm>

If this link does not work, please copy it to your browser and try again.  
If you want to print this pdf, we suggest you begin on the next page (2) to conserve paper.

**Devenez membre de la  
SOOA et soutenez la  
disponibilité future des pdfs  
gratuits sur ce site.**

<http://malimbus.free.fr/adhesion.htm>

Si ce lien ne fonctionne pas, veuillez le copier pour votre navigateur et réessayer.  
Si vous souhaitez imprimer ce pdf, nous vous suggérons de commencer par la page suivante  
(2) pour économiser du papier.

THE EGG COLOUR OF *Clamator levaillantii* (Swainson) IN NIGERIA - On 17 September 1975 a *Clamator levaillantii* trapped for ringing by R.E.S. was kept in a bird bag overnight when it laid an egg. It is characteristically broader and rounder than the eggs of the *Turdoides* species which are the usual hosts in Africa, and measures 25.3 x 19.5 mm. The breadth is 0.5 mm less than the smallest breadth of a series of ten *C. levaillantii* eggs recorded by Harrison (1971). The shell weight could not be determined because the egg was unblown and contained dried-up soft matter. The surface is smooth and slightly glossy and uniformly pale immaculate blue with a dingy grey tinge, the latter perhaps due to the dried-up contents.

*Clamator levaillantii* eggs in South and East Africa are pale blue to bluish-green, but in Nigeria eggs attributed to *C. levaillantii* on strong presumptive evidence are pink, matching the aberrant pink variety of *Turdoides plebejus* egg (the host species) sometimes found in Nigeria (Serle 1939). The egg here described shows that the blue-coloured *C. levaillantii* egg also occurs in Nigeria where *Turdoides plebejus* eggs show a remarkable variation in colour, from blue through grey-blue and mauve to pink.

R. E. Sharland and W. Serle

#### References

- Harrison, C. J. O. 1971 Notes on the identification of eggs, egg mimicry and distributional history and the status of the form *serratus*, in the parasitic *Clamator* cuckoos. Bull. Br. Orn. Cl. 91 : 126-131
- Serle, W. 1939 Field observations on some northern Nigerian birds Ibis 14(3) : 654-699

THE FOOD PASS MOVEMENT IN RAPTORS - This movement, characteristic of certain raptors, has been described by P. Brown (1969, Birds of Prey, A Survival Book, Andre Deutsch), quoting Balfour, as it is performed by the Hen Harrier *Circus cyaneus*. On two occasions at Enugu, I have seen what appeared to be a comparable manoeuvre, performed once by a pair of Black-shouldered Kites *Elanus caeruleus* and once by a pair of Shikras *Accipiter badius*. The birds passed one another at high speed on opposite but parallel upper and lower flight paths and it appeared that something was "passed" between them in the air.

S. G. Cowper

BLACK DUCK *Anas sparsa* ON MAMBILLA PLATEAU, FIRST RECORD FOR NIGERIA - During the period October 1974 to January 1975, two sightings were obtained of a duck in a small marsh near Yelwa on Mambilla Plateau, in October and December. On both occasions I had only brief glimpses but